



Coastal Wetland Invertebrate Communities

Formerly Indicator #4501

Overall Assessment

Status: Not Assessed

Trend: Not Assessed

Rationale: Part of an overall analysis of biological communities of Great Lakes coastal wetlands.

Note: This is a progress report towards implementation of this indicator. The indicator is currently being used throughout the entire Great Lakes basin, but data will not be available until 2012. The following evaluation was constructed using input from investigators collecting invertebrate community composition data from Great Lakes coastal wetlands over the last several years. Regarding the following, neither experimental design nor statistical rigor has been used to specifically address the status and trends of invertebrate communities of coastal wetlands of the five Great Lakes. However, in the spring of 2011, an effort was put forth by a consortium of universities that established a statistically sound basin-wide coastal wetland monitoring program. This indicator will be used, along with others, at the majority of coastal wetlands with a surface water connection to the Great Lakes that are greater than 4 hectares in size. The effort is bi-national and basin wide and will produce scientifically-defensible information on the status and trends of Great Lakes coastal wetlands.

Lake-by-lake Assessment

Each lake was categorized with a not assessed status and an undetermined trend, indicating that data were not available yet.

Purpose

- To directly measure specific components of invertebrate community composition
- To infer the chemical, physical and biological integrity and range of degradation of Great Lakes coastal wetlands

Ecosystem Objective

Significant wetland areas in the Great Lakes System that are threatened by urban and agricultural development and waste disposal activities should be identified, preserved and, where necessary, rehabilitated (Annex 13 GLWQA). Conducting monitoring and surveillance activities will gather definitive information on the location, severity, aerial or volume extent, and frequency of the Great Lakes coastal wetlands (Annex 11 GLWQA). This indicator supports the restoration and maintenance of the chemical, physical and biological integrity of the Great Lakes basin and beneficial uses dependent on healthy wetlands (Annex 2 GLWQA).

State of the Ecosystem

Teams of Canadian and American researchers from several research groups (e.g. the Great Lakes Coastal Wetlands Consortium, the Great Lakes Environmental Indicators project investigators, the U.S. Environmental Protection Agency (U.S. EPA) Regional Environmental Monitoring and Assessment Program (REMAP) group of researchers, and others) sampled large numbers of Great Lakes wetlands. In 2002 the Great Lakes Coastal Wetlands Consortium conducted extensive surveys of wetland invertebrates of the four lower Great Lakes. The Consortium-adopted Index of Biotic Integrity (IBI, Uzarski *et al.* 2004) was applied in wetlands of northern Lake Ontario. The results can be obtained from Environment Canada (Environment Canada and Central Lake Ontario Conservation Authority 2004).

These methods are now being used basin-wide by a consortium of universities but these data will not be available until 2012.

Uzarski *et al.* (2004) collected invertebrate data from 22 wetlands in Lake Michigan and Lake Huron during 1997 through 2001. They determined that wetland invertebrate communities of northern Lakes Michigan and Huron generally produced the highest IBI scores. IBI scores were primarily based on richness and abundance of Odonata, Crustacea plus Mollusca taxa richness, total genera richness, relative abundance Gastropoda, relative abundance Sphaeriidae, Ephemeroptera plus Trichoptera taxa richness, relative abundance Crustacea plus Mollusca, relative abundance Isopoda, Evenness, Shannon Diversity Index, and Simpson Index. Wetlands near Escanaba and Cedarville, Michigan, scored lower than most in the area. A single wetland near the mouth of the Pine River in Mackinac County, MI, consistently scored low. In general, all wetlands of Saginaw Bay scored lower than those of northern Lakes Michigan and Huron. However, impacts are more diluted near the outer bay and IBI scores reflect this. Wetlands near Quanicassee and Almeda Beach, MI, consistently scored lower than other Saginaw Bay sites.

Burton and Uzarski also studied drowned river mouth wetlands of eastern Lake Michigan quite extensively since 1998. Invertebrate communities of these systems show linear relationship with latitude. However, this relationship also reflects anthropogenic disturbance. Based on the metrics used (Odonata richness and abundance, Crustacea plus Mollusca richness, total genera richness, relative abundance Isopoda, Shannon Index, Simpson Index, Evenness, and relative abundance Ephemeroptera), the sites studied were placed in increasing community health in the order Kalamazoo, Pigeon, Muskegon, White, Pentwater, Pere Marquette, Manistee, Lincoln, and Betsie. The most impacted systems of eastern Lake Michigan are located along southern edge and impacts decrease to the north.

Wilcox *et al.* (2002) attempted to develop wetland IBIs for the upper Great Lakes using microinvertebrates. While they found attributes that showed promise during a single year, they concluded that natural water level changes were likely to alter communities and invalidate metrics. They found that Siskiwit Bay, Bark Bay, and Port Wing had the greatest overall taxa richness with large catches of cladocerans. They ranked microinvertebrate communities of Fish Creek and Hog Island lower than the other four western Lake Superior sites. Their work in eastern Lake Michigan testing potential metrics placed the sites studied in decreasing community health in the order Lincoln River, Betsie River, Arcadia Lake/Little Manistee River, Pentwater River, and Pere Marquette River. This order was primarily based on the median number of taxa, the median Cladocera genera richness, and also a macroinvertebrate metric (number of adult Trichoptera species).

Pressures

Physical alteration and eutrophication of wetland ecosystems continue to be a threat to invertebrates of Great Lakes coastal wetlands. Both can promote establishment of non-native vegetation, and physical alteration can destroy plant communities altogether while changing the natural hydrology to the system. Invertebrate community composition is directly related to vegetation type and densities; changing either of these components will negatively impact the invertebrate communities.

Agriculture

Agriculture degrades wetlands in several ways, including nutrient enrichment from fertilizers, increased sediments from erosion, increased rapid runoff from drainage ditches, introduction of agricultural non-native species (reed canary grass), destruction of inland wet meadow zone by plowing and diking, and addition of herbicides.

Urban development

Urban development degrades wetlands by hardening shoreline, filling wetland, adding a broad diversity of chemical

STATE OF THE GREAT LAKES 2012 - DRAFT

pollutants, increasing stream runoff, adding sediments, and increased nutrient loading from sewage treatment plants. In most urban settings, almost complete wetland loss has occurred along the shoreline.

Residential shoreline development

Along many coastal wetlands, residential development has altered wetlands by nutrient enrichment from fertilizers and septic systems, shoreline alterations for docks and boat slips, filling, and shoreline hardening. Agriculture and urban development are usually less intense than local physical alteration which often results in the introduction of non-native species.

Mechanical alteration of shoreline

Mechanical alteration takes a diversity of forms, including diking, ditching, dredging, filling, and shoreline hardening. With all of these alterations, non-native species are introduced by construction equipment or in introduced sediments.

Introduction of non-native species

Non-native species are introduced in many ways. Some were purposefully introduced as agricultural crops or ornamentals, later colonizing in native landscapes. Others came in as weeds in agricultural seed. Increased sediment and nutrient enrichment allow many of the worst aquatic weeds to out-compete native species. Most of the worst non-native species are either prolific seed producers or reproduce from fragments of root or rhizome. Non-native animals have also been responsible for increased degradation of coastal wetlands.

Pressures were described by Dennis Albert in the Coastal Wetland Plant Communities Indicator # 4862

Management Implementations

Although monitoring protocols have been developed for this indicator by the Great Lakes Coastal Wetlands Consortium, monitoring on basin wide scale has not yet occurred. Implementations of a long term coastal wetland monitoring program is pending, however support for this program is needed by resource managers throughout the basin.

Assessing Data Quality

Insert “x” under the statement that best corresponds with each data characteristic

Data Characteristics	Strongly Agree	Agree	Neutral or Unknown	Disagree	Strongly Disagree	Not Applicable
1. Data are documented, validated, or quality-assured by a recognized agency or organization	X					
2. Data are traceable to original sources	X					
3. The source of the data is a known, reliable and respected generator of data	X					
4. Geographic coverage and scale of data are appropriate to the Great Lakes basin	X					
5. Data obtained from sources within the U.S. are comparable to those from Canada	X					

STATE OF THE GREAT LAKES 2012 - DRAFT

6. Uncertainty and variability in the data are documented and within acceptable limits for this indicator report	X					
--	---	--	--	--	--	--

Clarifying Notes:

Acknowledgments

Authors:

Donald G. Uzarski, Institute for Great Lakes Research, CMU Biological Station, and Department of Biology, Central Michigan University, Mount Pleasant, MI.

Thomas M. Burton, Departments of Zoology and Fisheries and Wildlife, Michigan State University, East Lansing, MI, 48824. (2006)

Contributors:

Danielle J. Sass, Oak Ridge Institute of Science and Education (ORISE) Research Fellow, Appointed to the U.S. Environmental Protection Agency (U.S. EPA), Great Lakes National Program Office (GLNPO) (2008)

Sources

Environment Canada and Central Lake Ontario Conservation Authority. 2004. *Durham Region Coastal Wetland Monitoring Project: year 2 technical report*. Downsview, ON. ECB-OR.

Uzarski, D.G., Burton, T.M., and Genet, J.A. 2004. Validation and performance of an invertebrate index of biotic integrity for Lakes Huron and Michigan fringing wetlands during a period of lake level decline. *Aquat. Ecosystem Health & Manage.* 7(2):269-288.

Wilcox, D.A., Meeker, J.E., Hudson, P.L., Armitage, B.J., Black, M.G., and Uzarski, D.G. 2002. Hydrologic variability and the application of index of biotic integrity metrics to wetlands: a Great Lakes evaluation. *Wetlands* 22(3):588-615

Last Updated

State of the Great Lakes 2009 report.
An editor’s note was added for the 2011 reporting cycle.